

**Measure Information Form**  
**Collected For: CMS Outcome Measures (Claims Based)**

**Measure Set:** CMS Mortality Measures

**Set Measure ID#:** MORT-30-STR

**Performance Measure Name:** Hospital 30-day, all-cause, risk-standardized mortality rate following acute ischemic stroke hospitalization.

**Description:** The measure estimates a hospital-level, risk-standardized mortality rate for patients discharged from the hospital with a principal diagnosis of acute ischemic stroke.

**Rationale:** Risk-standardized mortality rates (RSMRs) can provide important additional information about quality of care that is not currently captured by the process measures and is currently unavailable to hospitals. Variation in mortality, after adjusting for case-mix, may reflect differences in hospitals' general environments (such as coordination of care, patient safety policies, and staffing) or variation in care processes. Outcome measures can focus attention on a broad set of healthcare activities that affect patients' well-being. Moreover, improving outcomes is the ultimate goal of quality improvement, and so the inclusion of outcomes measures assists in attaining improvement goals.

Stroke is a common condition with substantial mortality and morbidity. The condition imposes a substantial burden on patients and the health care system, and there is marked variation in outcomes by institution.

**Type of Measure:** Outcome

**Improvement Noted As:** A decrease in the RSMR.

**Numerator Statement:**

This outcome measure does not have a traditional numerator and denominator like a core process measure (e.g., percentage of adult patients with diabetes aged 18-75 years receiving one or more hemoglobin A1c tests per year); thus, we are using this field to define our outcome. The calculation of the rate is defined below under Measure Calculation.

The outcome for this measure is 30-day all-cause mortality. We define mortality as death from any cause within 30 days after the index admission date.

**Denominator Statement:**

The target population for this measure includes admissions for Medicare Fee-for-Service (FFS) beneficiaries aged ≥65 years discharged from acute care non-federal hospitals, having a principal discharge diagnosis of acute ischemic stroke.

**Included Populations:** Admissions for Medicare FFS beneficiaries aged ≥65 years discharged from non-federal acute care hospitals, having a principal discharge diagnosis of acute ischemic stroke.

CMS FFS beneficiaries hospitalized within an acute care non-federal hospital are included if they have been enrolled in Part A and Part B Medicare for the 12 months prior to the date of admission to ensure a full year of administrative data for risk-adjustment.

For patients with more than one admission in a given year for a given condition, only one admission is randomly selected to include in the cohort (others are excluded).

The measure includes patients who are admitted to an acute care hospital with a diagnosis of stroke and then transferred to another acute facility if the primary discharge diagnosis is stroke at the second hospital. The measure considers admission to the first hospital as the start of an acute episode of care and assigns the patient's outcome to the hospital that initially admitted them.

#### **ICD-9-CM codes that define the patient cohort**

433.01	Occlusion and stenosis of precerebral arteries, Basilar artery with cerebral infarction
433.11	Occlusion and stenosis of precerebral arteries, Carotid artery with cerebral infarction
433.21	Occlusion and stenosis of precerebral arteries, Vertebral artery with cerebral infarction
433.31	Occlusion and stenosis of precerebral arteries, Multiple and bilateral with cerebral infarction
433.81	Occlusion and stenosis of precerebral arteries, Other specified precerebral artery with cerebral infarction
433.91	Occlusion and stenosis of precerebral arteries, Unspecified precerebral artery with cerebral infarction, Precerebral artery NOS
434.01	Occlusion of cerebral arteries, Cerebral thrombosis with cerebral infarction, Thrombosis of cerebral arteries
434.11	Occlusion of cerebral arteries, Cerebral embolism with cerebral infarction
434.91	Occlusion of cerebral arteries, Cerebral artery occlusion, unspecified, with cerebral infarction

#### **Excluded Populations:**

The measure excludes admissions for patients:

- with inconsistent or unknown mortality status or other unreliable data (e.g. date of death precedes admission date)
- who were transferred from another acute care hospital (because the death is attributed to the hospital where the patient was initially admitted)
- enrolled in Medicare Hospice programs any time in the 12 months prior to the index hospitalization including the first day of the index admission (since it is likely these patients are continuing to seek comfort measures only)
- who were discharged against medical advice (AMA) (because providers did not have the opportunity to deliver full care and prepare the patient for discharge)

**Risk Adjustment:** For each patient, covariates are obtained from administrative data extending 12 months prior to, and including, the index admission. For all patients, information from Medicare inpatient claims, physician Part B claims and hospital outpatient claims are used for risk adjustment. Inpatient claim records have data on hospitalization for and include demographic information, principal and secondary diagnosis codes, and procedure codes. Diagnosis codes for comorbidities are also collected from physician and hospital outpatient files. These data are captured from the claim(s) for the index admission and from all inpatient and outpatient claims for the entire year before the patient's index stroke hospitalization to be utilized in the risk-adjustment model. This measure also includes risk-adjustment for patients that are seen at an outside Emergency Department (ED) and then transferred to a hospital for their index admission.

Only variables that convey information about patients' clinical status at the time of admission are used for the risk-adjustment, while complications that arise during the course of patients' index hospitalization are not included in the model.

Full details of the development of the risk-standardization model for this measure are available at: <http://www.qualitynet.org>.

The final set of risk-adjustment variables included:

Demographics	Age Male
Cardiovascular/ Cerebrovascular	Congestive Heart Failure Valvular and Rheumatic Heart Disease Congenital Cardiac/Circulatory Defects Hypertensive Heart Disease Specified Heart Arrhythmias Cerebral Hemorrhage Ischemic or Unspecified Stroke Precerebral Arterial Occlusion and Transient Cerebral Ischemia Cerebral Atherosclerosis and Aneurysm Hemiplegia/Hemiparesis History of Infection Metastatic Cancer and Acute Leukemia and Other Major Cancers Lymphatic, Head and Neck, Brain, Breast, Colorectal and Other Major Cancers Protein-Calorie Malnutrition Other Significant Endocrine and Metabolic Disorders Other Gastrointestinal Disorders Disorders of the Vertebrae and Spinal Discs Osteoarthritis of Hip or Knee Other Musculoskeletal and Connective Tissue Disorders Iron Deficiency and Other/Unspecified Anemia and Blood Disease Dementia or senility Major Psychiatric Disorders Quadriplegia, Other Extensive Paralysis Multiple Sclerosis/Other Neurological Conditions

	Seizure Disorders and Convulsions Hypertension Peripheral Vascular Disease Chronic Obstructive Pulmonary Disease Pneumonia Pleural Effusion/Pneumothorax Other Eye Disorders Other Ear, Nose, Throat, and Mouth Disorders Dialysis Status Renal Failure Urinary Tract Infection Male Genital Disorders Decubitus Ulcer of Skin Chronic Ulcer of Skin, Except Decubitus Other Dermatological Disorders
Other	ED-transfer

**Model Validation:** Hospital-specific risk-standardized mortality estimates derived from this claims-based model were compared to hospital-specific RSMRs based on a model developed using medical record data including a measure of stroke severity from the Medicare Health Care Quality Improvement Program's National Stroke Project (NSP). The correlation coefficient of the RSMRs from the claims-based and medical record models was 0.80.

**Data Accuracy:** The administrative claims data used to calculate the measure are maintained by CMS' Office of Information Services. These data undergo additional quality assurance checks during measure development and maintenance.

**Measure Analysis Suggestions:** None

**Sampling:** No.

**Data Reported As:** Hospital 30-day, all-cause, risk-standardized mortality rate (RSMR) following acute ischemic stroke hospitalization.

**Measure Calculation:**

The measure estimates hospital-level 30-day all-cause RSMR for stroke using hierarchical logistic regression modeling. In brief, the approach simultaneously models two levels (patient and hospital) to account for the variance in patient outcomes within and between hospitals. At the patient level, the model adjusts the log-odds of mortality within 30 days of admission for age, sex, selected clinical covariates, and a hospital-specific intercept. At the hospital level, it models the hospital-specific intercepts as arising from a normal distribution. The hospital intercept represents the underlying risk of mortality at the hospital, after accounting for patient risk. The hospital-specific intercepts are given a distribution in order to account for the clustering (non-independence) of patients within the same hospital. If there were no differences among hospitals, then after adjusting for patient risk, the hospital intercepts should be identical across all hospitals.

The RSMR is calculated as the ratio of the number of “predicted” deaths to the number of “expected” deaths, multiplied by the national unadjusted mortality rate. For each hospital, the “numerator” of the ratio is the number of deaths within 30 days predicted on the basis of the hospital’s performance with its observed case-mix, and the “denominator” is the number of deaths expected on the basis of the nation’s performance with that hospital’s case-mix. This approach is analogous to a ratio of “observed” to “expected” used in other types of statistical analyses. It conceptually allows for a comparison of a particular hospital’s performance given its case-mix to an average hospital’s performance with the same case-mix. Thus, a lower ratio indicates lower-than-expected mortality or better quality, and a higher ratio indicates higher-than-expected mortality or worse quality.

The predicted number of deaths (the numerator) is calculated by regressing the risk factors and the hospital-specific intercept on the risk of mortality, multiplying the estimated regression coefficients by the patient characteristics in the hospital, transforming, and then summing over all patients attributed to the hospital to get a value. The expected number of deaths (the denominator) is obtained by regressing the risk factors and a common intercept on the mortality outcome using all hospitals in our sample, multiplying the subsequent estimated regression coefficients by the patient characteristics observed in the hospital, transforming, and then summing over all patients in the hospital to get a value. To assess hospital performance in any reporting period, we re-estimate the model coefficients using the years of data in that period.

The statistical modeling approach is described fully in the original methodology report.

### **Selected References:**

- Bernheim S, Wang C, Wang Y, et al. 2010. Hospital 30-Day Mortality Following Acute Ischemic Stroke Hospitalization Measure: Methodology Report.
- Bernheim S, Wang C, Wang Y. 2013. 2013 Measure Updates and Specifications Report: Hospital 30-day Mortality Following an Admission for an Acute Ischemic Stroke (Version 2.0).
- Bradley EH, Herrin J, Elbel B, et al. Hospital Quality for Acute Myocardial Infarction: Correlation Among Process Measures and Relationship With Short-term Mortality. *The Journal of the American Medical Association* 2006;296:72-8.
- Bravata DM, Ho SY, Meehan TP, Brass LM, Concato J. Readmission and death after hospitalization for acute ischemic stroke: 5-year follow-up in the Medicare population. *Stroke* 2007;38:1899-904.
- Bravata DM, Wells CK, Lo AC, et al. Processes of Care Associated With Acute Stroke Outcomes. *Archives of Internal Medicine* 2010;170:804-10.
- Fonarow GC, Reeves MJ, Zhao X, et al. Age-Related Differences in Characteristics, Performance Measures, Treatment Trends, and Outcomes in Patients With Ischemic Stroke. *Journal of the American Heart Association* 2010;121:879-91.
- Krumholz HM, Brindis RG, Brush JE, et al. 2006. Standards for Statistical Models Used for Public Reporting of Health Outcomes: An American Heart Association Scientific Statement From the Quality of Care and Outcomes Research Interdisciplinary Writing Group: Cosponsored by the Council on Epidemiology and Prevention and the Stroke Council Endorsed by the American College of Cardiology Foundation. *Circulation* 113: 456-462.
- Normand S-LT, Shahian DM. 2007. Statistical and clinical aspects of hospital outcomes profiling. *Stat Sci* 22(2):206-226.
- Saposnik G, Hill MD, O'Donnell M, Fang J, Hachinski V, Kapral MK. Variables Associated With 7-Day, 30-Day, and 1-Year Fatality After Ischemic Stroke. *Journal of the American Heart Association* 2008;39.